

Synonym

CD70,CD27LG,TNFSF7,TNFSF7G,CD27L

Source

Human CD27 Ligand, Llama IgG2b Fc Tag(CDL-H5255) is expressed from human 293 cells (HEK293). It contains AA Ser 52 - Pro 193 (Accession # [NP_001243.1](#)).

Predicted N-terminus: Glu

Molecular Characterization

This protein carries a llama IgG2b Fc tag at the N-terminus

The protein has a calculated MW of 75.5 kDa. The protein migrates as 140-160 kDa under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.01 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE

Human CD27 Ligand, Llama IgG2b Fc Tag on SDS-PAGE under non-reducing (NR) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA



Immobilized Human CD27 Ligand, Llama IgG2b Fc Tag (Cat. No. CDL-H5255) at 2 µg/mL (100 µL/well) can bind Human CD27, His Tag (Cat. No. CD7-H522b) with a linear range of 4-63 ng/mL (QC tested).



Immobilized Human CD27 Ligand, Llama IgG2b Fc Tag (Cat. No. CDL-H5255) at 2 µg/mL (100 µL/well) can bind Biotinylated Human CD27, Fc, Avitag (Cat. No. TN7-H82F6) with a linear range of 0.3-20 ng/mL (Routinely tested).

Background

Cluster of Differentiation 70 (CD70) is also known as CD27 ligand (CD27L / CD27LG), TNFSF7, TNFSF7G, is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and is a surface antigen found on activated T-and B-lymphocytes and mature dendritic cells. Binding of CD70 to its receptor CD27 induces in priming, effector functions, differentiation and memory formation of T-cells, and thus is involved in the biological processes including T-cell activation, the proliferation of costimulated T-cells, as well as the generation of cytolytic T-cells. CD70 on T cells provides costimulatory signals that are required for T cell proliferation, clonal expansion and the promotion of effector T cell formation. CD70 on mouse B cell has been shown to inhibit terminal differentiation of activated B cells into plasma cells and enhances commitment to memory B cell responses. CD70 induces proliferation and IFN γ production, on NK cells.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.