



Synonym

HDNF, Nerve growth factor 2, NGF-2, Neurotrophic factor, NTF3, NT3, Neurotrophin3

Source

Human NT-3 Protein, Tag Free (NT3-H5213) is expressed from human 293 cells (HEK293). It contains AA Tyr 139 - Thr 257 (Accession # [P20783](#)).
Predicted N-terminus: Tyr 139

Molecular Characterization

NT-3(Tyr 139 - Thr 257)
P20783

This protein carries no "tag".

The protein has a calculated MW of 13.6 kDa. The protein migrates as 13-14 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.01 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>95% as determined by SEC-HPLC.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human NT-3 Protein, Tag Free on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-HPLC




The purity of Human NT-3 Protein, Tag Free (Cat. No. NT3-H5213) was greater than 95% as determined by SEC-HPLC.

Bioactivity-ELISA

Discounts, Gifts,
and more!





Immobilized Human NT-3 Protein, Tag Free (Cat. No. NT3-H5213) at 1 µg/mL (100 µL/well) can bind Human TrkC Protein, Fc Tag (Cat. No. TRC-H5256) with a linear range of 0.8-13 ng/mL (QC tested).

Background

Neurotrophin-3 (NT-3), encoded by the NTF3 gene, is a member of the NGF family of neurotrophic factors (also named neurotrophins). The neurotrophin family is comprised of at least four proteins including NGF, BDNF, NT-3, and NT-4/5. These secreted cytokini.

Clinical and Translational Updates

Discounts, Gifts,
and more!

